

12

EUROPEAN PATENT APPLICATION

21 Application number: 88307636.6

51 Int. Cl.4: **A 61 K 31/40**

22 Date of filing: 18.08.88

30 Priority: 20.08.87 US 87432

43 Date of publication of application:
22.02.89 Bulletin 89/08

84 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

71 Applicant: **MONOCLONETICS INTERNATIONAL, INC.**
18333 Egret Bay Boulevard Suite 270
Houston Texas 77058 (US)

72 Inventor: **De La Parra, Jacinto Juarez**
122 Blond Petersen
Ermita Prado Mexico City MX DF (MX)

74 Representative: **Bizley, Richard Edward et al**
BOULT, WADE & TENNANT 27 Farnival Street
London EC4A 1PQ (GB)

54 **Taliscanin and other Aristolactams for treating neurological disorders, Parkinson's disease, Alzheimer disease and impotence.**

57 Aristolactams e.g. Taliscanin may be used in preparing medication for treating neurological disorders, in particular Parkinson's disease, or impotence. The method involves administering an aristolactam, especially Taliscanin, an extract of the root of Aristolochia Taliscana, to an afflicted patient with a dosage varied in accordance with the patient's response, which dosage generally ranges from 280 to 4000 micrograms per day.

Description

**TALISCANIN AND OTHER ARISTOLACTAMS FOR TREATING NEUROLOGICAL DISORDERS,
PARKINSON'S DISEASE, ALZHEIMER DISEASE AND IMPOTENCE**

5 The invention relates to the use of Taliscanin, and other aristolactams, for preparing medication in the treatment of neurological disorders, Parkinson's disease, Alzheimer disease, and impotence.

Among neurological disorders, Parkinson's disease is one of the most common. In the United States alone, there are over one million patients who suffer from it. In addition, the number of Parkinson patients is constantly increasing. The increase is because the disease is an illness of the middle and late years of life, and people are today living longer. Further, the patients with the disease tend to live on for many years, hence, their numbers accumulate.

10 Parkinson's disease most frequently begins between the ages of forty and seventy, with a peak age of onset being in the sixth decade. It is estimated that about one percent of the population over the age of fifty is afflicted.

15 The characteristic symptoms of Parkinson's disease include movement problems, such as impedence of alternating movements, bradykinesia (slowness of movement), akinesia, or hypokinesia, a tremor (which occurs at rest but not usually during voluntary movement), stooped posture, rigidity, and an expressionless face.

20 The onset of the disease is characterized by a reduction in the rate of blinking, relative immobility and poverty of movement, and the characteristic tremor. As the disease develops the poverty of movement becomes more apparent, and is shown by infrequency of swallowing, slowness of chewing, disinclination to adjust the position of the body and limbs, and the lack of movements of cooperation, for example, arising from a chair without first adjusting the feet.

Patients frequently exhibit a festinating gait -- losing balance and "chasing" the center of gravity to avoid losing balance. Rigidity (alternating or continuous firming and tensing of the muscles), usually also appears. Eventually, due to tensing of the frontal muscles, the patient takes on a stooped posture. Similarly, tensing of the flexor muscles in the hands and feet can cause a permanent bent attitude of these extremities.

25 Potentially the most serious problem for the patient is dementia, which is commonly associated with Parkinson's disease. In one study it was reported that the incidence of dementia in Parkinson's disease patients was ten times as high as for their age matched spouses. Parkinson's patients with dementia have been reported to become more seriously involved in a shorter period of time and to respond less well to standard L-dopa therapy.

30 Parkinson's disease can be of three separate types: the post-encephalitic, idiopathic, and the arteriosclerotic type. The post-encephalitic type is not true Parkinson's disease inasmuch as it is not degenerative. It stems from an infection that involves not only the basal ganglia but the entire nervous system. The infection was caused by a virus which appeared during the worldwide influenza epidemics of 1917 to 1927.

35 The arteriosclerotic type, contrary to what was earlier believed, is now thought to be the same as the idiopathic ("unknown cause") type. The same symptoms occur, except that they are accompanied by arteriosclerosis. The arteriosclerotic type usually occurs in older people and involves more severe symptoms than the more common idiopathic type.

40 Physiologically speaking, Parkinsonism is a disease of the extrapyramidal motor system. This system governs automatic, static, postural, and other motor activities of the nervous system which are not readily modifiable. On the anatomical level, however, Parkinsonism is a progressive disease of the basal ganglia, i.e., the small center or nucleus at the base of the brain.

45 The extrapyramidal motor system is believed to be controlled by the basal ganglia and certain other brainstem nuclei. The striatum and the substantia nigra are both component structures of the basal ganglia. Parkinsonism is believed to be caused by aberrations in the neurotransmitters in the striatum and the substantia nigra, which in turn cause degeneration of neurons in those areas.

Neurotransmitters are substances which are synthesized and stored in the presynaptic terminals. A synapse is a functional junction between two neurons, where a nerve impulse is transmitted from one neuron to another. In response to an appropriate stimulus, a neurotransmitter is released across the synaptic gap to combine with specific receptor sites on the postsynaptic neuron.

50 The most important neurotransmitters from the point of view of basal ganglionic function are acetylcholine, dopamine, gamma aminobutyric acid (GABA), and serotonin. Acetylcholine exists at high concentrations in the striatum. It is synthesized and released by the small (Golgi type 2) neostriatal neurons, upon which it has an excitatory effect.

55 Dopamine is one of the family of neurotransmitters known as catecholamines, the others in that family being epinephrine, and norepinephrine. Dopamine is a precursor in the synthesis of epinephrine, which in turn is a precursor for norepinephrine. The areas of the brain which are richest in dopamine are the substantia nigra and the striatum, where it is localized in synaptic endings of nigral fibers. Stimulation of the substantia nigra causes a release of dopamine, which has an inhibitory effect on the neostriatal neurons.

60 Normally, the extrapyramidal system functions properly due to the interaction between two antagonistic systems. One of these systems is to some extent regulated by serotonin and the catecholamines, and the other by acetylcholine and histamine. The interaction between these two antagonistic systems is known as the

balance between the wakeful and sleep states, or the circadian cycle.

In Parkinson's disease the levels of the catecholamines are reduced. In particular, concentrations of dopamine are greatly decreased in the striatum and the substantia nigra. This may result either because the synthesis of dopamine is blocked, or because the enzymes which metabolize it (monoamine oxidase) are increased.

The decreased release of dopamine in the striatum which occurs in Parkinson's disease disinhibits the neurons that synthesize acetylcholine. This causes a predominance of cholinergic (acetylcholine-like) activity.

The most notable aspect of the pathology of Parkinson's disease is that there is a loss of pigmented cells in the substantia nigra and other pigmented nuclei. The normal ranges for nigral cells is about 425,000 for a young adult diminishing to about 200,000 at age eighty. In most Parkinson's patients there are less than 100,000. The degree of dopamine deficiency in the substantia nigra correlates with the degree of cell loss in this area.

Inasmuch as cholinergic activity predominates in Parkinsonism, one method of treatment is to administer anticholinergic drugs to restore the ratio between dopamine and acetylcholine. Typical anticholinergic drugs are belladonna and trihexyphenidyl. A related method of treatment involves administration of dopaminergic agents, e.g., L-dopa, amantadine, and bromocriptine. This causes an increase in the otherwise reduced levels of dopamine.

The problem with anticholinergic substances is that they cause a variety of undesirable side effects, including myocardium infarction, glaucoma, toxic psychosis, anxiety, hallucinations, nausea, vomiting, urine retention in prostatic hypertrophy, constipation, miosis (contraction of the pupil), intraocular hypertension, mouth dryness, dyschezias, and orthostatic hypotension. Dopaminergic agents cause other side effects, such as the induction of involuntary movements including restlessness, grimacing, choreoathetosis and dystonia of the limbs, neck, and trunk.

It is clear, therefore, that a drug which has less severe side effects than the dopaminergic and anticholinergic agents currently in use would be of great benefit in the treatment of Parkinsonism.

It is also suspected that other neurological disorders of the extrapyramidal system are caused by the same mechanism, and the same areas of the brain are affected, as in Parkinsonism. A number of other disorders can cause some or all of the symptoms of Parkinson's disease. For example, Alzheimer disease can cause elements of parkinsonian akinesia, rigidity, poor balance, and tremor. A drug which treats Parkinsonism should, therefore, all be useful in treatment of such neurological disorders.

The compound Taliscanin has, until now, been used in treating snake bites. It has never been used or suggested for use in treating neurological disorders, Alzheimer disease, or impotency.


It has been discovered that Taliscanin, an extract from the root of *Aristolochia Taliscana*, alleviates the symptoms of Parkinsonism and related neurological disorders. Taliscanin has also been found effective in the treatment of Alzheimer disease and impotency. Taliscanin should also be effective in treating paralysis which is secondary to cerebral vascular accidents or strokes, diabetes insipidus, choreoathetosis, obesity secondary to hypothalamic disorders and amenorrhea secondary to hypothalamic disorders, deafness of neurological origin and mutism.

The invention provides the use of an aristolactam in the preparation of a medicament for treating neurological disorders or impotence.


The invention further includes a pharmaceutical or veterinary formulation comprising an aristolactam, preferably Taliscanin, formulated for pharmaceutical or veterinary use in the treatment of neurological disorders or impotence, optionally in unit dosage form and/or optionally including an acceptable diluent, carrier or excipient.

Taliscanin has far fewer side effects than encountered with the conventional Parkinson's treatments such as dopaminergic and anticholinergic agents. Further, it is believed to actually regenerate at least some of the damaged or injured neurons of the substantia nigra, i.e., those which are not in the latter stages and, thus, may be effective in reducing aging and increasing longevity.

Taliscanin has the following chemical formula:



Taliscanin is one of the family known as aristolactams, which family has the following generic formula:



It is believed that several aristolactam derivatives may have the same therapeutic activity as does Taliscanin. Among these are the aristolams shown in Channer, "The Aristolochic Acids and Aristolactams," Isoquinoline Alkaloid Research, Plenum Press, Chapter 17 (1978), for example, aristolactam, aristored, aristolochic acid-D methyl ether lactam, aristolactam Beta-D- glucoside, aristolactam-A11, aristolactam-A111, aristolactam-B11 (cepharanone-B), aristolactam-B111, cepharanone-A, and doryflavine.

Taliscanin, and the aristolactams, are well-known compounds which are described in D.B. Mix et al. "The Aristolochic Acids and aristolactams," Journal of Natural Products, Vol. 45 No. 5 (1982). Taliscanin can be extracted from the root of Aristolochia Taliscana. The extraction process is well-known in the art, and is described in L. Maldonado et al., "La Taliscanina, un Componente de Aristolochia Taliscana" Cincia Mexico Vol. 24 at p. 237 (1966); Channer, "The Aristolochic Acids and Aristolactams," Isoquinoline Alkaloid Research, Plenum Press, Chapter 17 (1978). It should also be possible to synthesize Taliscanin, although that has not yet been done.

For the sake of completeness, one example of an extraction process follows.

Example 1: Extraction Process

9.1 kg of the root of Aristolochia Taliscana was pulverized and then suspended in undiluted hexane. The suspension was then acted on by a Soxhlet extractor until all the hexane was exhausted. The residue was then suspended in benzene, which was again exhausted by a Soxhlet extractor. 143.0 grams of the resulting brown residue was then dissolved in a 2:3 mixture of benzene-hexane, and the solution was fractionated through an alumina chromatography column. Polar fractions were then eluted with a 7:3 benzene-ether mixture.

This procedure yielded 13.8 grams of Taliscanin, having a melting point of 272-273°C, maxima UV absorption of 242 254, 310, 395 millimicrons; E of 38900, 35490, and 16220. It showed infra red absorption at 3450 cm^{-1} (CHCH₃), and 1690 cm^{-1} (NH) (carbenito of lactam) and a 1615 cm^{-1} aromatic double bond.

Example 2: Preparation of Medication

Clinical trials were conducted with Taliscanin to test its efficacy in treating Parkinsonism and other neurological disorders. The Taliscanin was prepared for oral administration to the patients by the following method.

100 ml of 96 proof ethanol was mixed with a sufficient amount of distilled water to bring the final alcohol solution to 60 proof. 20 mg of Taliscanin were added, and the solution was shaken.

Another batch was made up for particularly sensitive patients, the only difference being that in this solution the final alcohol concentration was 30 proof.

The alcohol/water solutions prepared were protected from sunlight by storage in amber glass. Storage can be at room temperature or under refrigerated conditions.

It should be emphasized that an alcohol/water solution is only one example of a suitable vehicle for Taliscanin. Other possible vehicles are distilled water, lactose, and dextrose/water, e.g., Dextrose Water 5.

Example 3: Administration of the Medication

The Taliscanin in solution was administered to the patients in the clinical trial at the rate of 280 micrograms of Taliscanin (three drops of solution) per day. The patients were checked once per week, or less often if weekly check-ups were too inconvenient.

The check-up involved monitoring the patient's rigid marche, facie, tremor, and supination pronation (forearm movement) for improvement. The degree of dilation of the pupils is an important parameter in the evaluation of the patient, and this was also monitored to determine whether it was normalizing. It was also determined whether the patient suffered from insomnia, which is a side effect of the drug where the dosage is too large. The only other known side effect is a painful erection in some males.

Patients that showed an improvement were kept on the same dosage of Taliscanin. For those that did not, the dosage was increased to 600 micrograms (six drops) per day, and they were followed up two weeks later. For any which still showed no improvement, the dosage was increased another three drops per day until there was an improvement or until the maximum dose was reached.

The maximum clinical dose was arrived at following a toxicity study on dogs. This study showed a lethal dose

of Taliscanin to be 4.0 mg for a 35 kg dog. Based on this result, it was estimated that 8.0 mg would kill a 70 kg man. The maximum clinical dose administered was reduced considerably from the 8.0 mg level. It was, in fact, only 4.0 mg per day.

The medication is taken orally every one to six hours as required to administer the daily dosage. To prevent insomnia in patients receiving a total of 840 micrograms per day or more, the medication should not be given later than two hours before bedtime.

It is recommended that 1 ml per week of Hyaluronidase (Wydase T.M.) be administered intramuscularly for the first four to six months of treatment. Hyaluronidase increases the permeability to Taliscanin of the blood/brain barrier and of the pre and postsynaptic terminals. However, Hyaluronidase is contraindicated for patients who are pregnant or who tend to bleed, for example, due to hemophilia.

Other agents which increase the permeability of the blood/brain barrier and/or the pre and postsynaptic terminals, for example Papavaina (available in Mexico), can be substituted for Hyaluronidase.

Once the symptoms of Parkinsonism, e.g., tremor and impaired motion, are no longer apparent the treatment can be stopped. To withdraw from medication, the dosage is gradually tapered off in increments of three drops per week. Gradual withdrawal is recommended because a sudden cessation of medication can, in some instances, cause relapse. Although relapse is not a common problem, caution nevertheless suggests a gradual withdrawal.

An alternative to complete cessation of medication is placing the patient on a maintenance dosage. This course is recommended where symptoms reappear as the dosage is decreased. The maintenance dosage is usually from 50 to 100 micrograms per day.

As mentioned above, Taliscanin was also effective in treating impotence and Alzheimer disease. The normal dosage for the treatment of impotence is 280 to 840 micrograms per day, and for Alzheimer Disease, it is 280 to 1200 micrograms per day.

It is emphasized that the medication can be administered by a number of methods other than orally, e.g., intravenously or intramuscularly. It can also be given at a variety of dosages. The parameters listed above are merely those that were used in these clinical trials.

Clinical Results

Taliscanin was administered to several thousand patients with various neurological disorders and Parkinsonism symptoms, using the above-described protocol. The results of the treatment, as broken down by patients showing predominant symptoms, appear in Table I below. Of these patients, approximately one-third of those in the first two (early) stages of Parkinson's disease were completely cured. Patients in the third and fourth (late) stages of the disease were not cured but were, in most cases, improved by Taliscanin.

Taliscanin was also effective in curing and improving various other neurological disorders related to and/or causing the same symptoms as Parkinsonism. See Table II below.

TABLE 1
EXTRAPYRAMIDAL SYSTEM SYMPTOMS (CLINICAL CASES) INDICATING
DESCENDING NERVE TRACTS THAT DO NOT ENTER INTO THE
FORMATION OF THE PYRAMIDS OF THE MEDULLA

Symptoms	Number of Patients				Success* Rate
	Total	Cure	Moderate Improvement	Without Improvement	
HYPERKINESIA (rapid movements)	2,236	652	1,390	194	91.32%
AKINESIA (low motility)	2,603	923	1,532	148	94.31%
HYPERTONIA (excess tension of muscles or arteries)	1,415	625	706	84	94.06%
HYPOTONIA (reduced tension)	375	54	248	73	80.53%
AUTONOMIC DYSFUNCTION (dysfunction of independent organs)	157	70	71	16	90.00%
RESTING TREMOR - SYNDROME OF PARKINSON (trembling in a static pose)	378	168	192	18	95.24%
INTENTION OR ACTION TREMOR AND TREMOR WITH A LESION IN THE CEREBELLUM (tremor in a moving state; active, static)	173	39	108	26	85.00%
CHOREA-ATHETOSIS (spasmodic movement of facial and extremity muscles)					
CHOREA, HEMIBALLISM, DYSTONIA, ATHETOSIS (involuntary movements on one side of the body; state of hyper- tonicity or hypotonicity)	44	21	22	1	97.73%
PROPULSION MARCHE (walking too fast and falling)	142	90	43	9	93.66%
OPTOKINETIC INSTAGMUS (eye twitching)	36	8	22	6	83.33%
DYSSYNERGIA (loss of power of muscular coordination)	504	166	306	32	93.65%

*Number of patients cured together with those showing moderate improvement as a percentage of the total.

side of the body; state of hypertonicity of hypotonicity) 44|21|22|1

TABLE II
TREATMENT WITH TALISCANIN OF PARKINSON'S DISEASE AND RELATED
NEUROLOGICAL DISORDERS

Etiology	Total	Cure	Number of Patients		
			Moderate Improvement	Mild Improvement	Without Improvement
VIRAL INFECTIONS:					
Encephalitis lethargica (viral)	596	229	244	56	67
Influenza	633	299	209	65	60
SYPHILIS	105	24	44	18	19
TUBERCULOSIS	766	270	300	100	96
VICERAL MYCOSIS (fungal)	146	33	43	23	47
CYSTICERCOSIS (parasitic)	54	38	37	8	9
EXOGENOUS INTOXICATION BY CARBON MONOXIDE	50	in majority of patients	5	4	3
EXOGENOUS INTOXICATION BY MANGANESE	45	-	23	9	13
COLLATERAL EFFECTS OF DRUGS:					
Antipsychotics:					
a) Largactil	324	48	140	78	58
b) Sinogan	32	7	6	15	4
c) Piportil	88	32	33	11	12
d) Haldol	90	28	35	15	12
Blood Pressure					
a) Reserpina	436	121	172	54	89
HEAD TRAUMA	59	29	13	7	10
CEREBRALVASCULAR SCLEROSIS	375	165	149	38	23
ATHEROSCLEROSIS (Parkinson's Disease)	89	32	36	18	3
IDIOPATHIC PARKINSON'S DISEASE	2,155	497	767	439	452
SYNDROMES:					
a) Von Monakov (viral)	61	36	11	4	10
b) Delhermitte van Bogaert	130	48	44	12	26
c) De Parinaud	31	12	13	5	1

It can be seen that Taliscanin was effective in curing or improving the vast majority of patients with Parkinsonian

It can be seen that Taliscanin was effective in curing or improving the vast majority of patients with Parkinsonian symptoms, Parkinson's disease, and other neurological disorders. Because of the similarity in structural formula, it is believed that other aristolactams will also be effective in treating these disorders. It should be understood that all descriptions herein are exemplary only.

Claims

1. The invention provides the use of an aristolactam in the preparation of a medicament for treating neurological disorders or impotence.

2. The use of claim 1 wherein the neurological disorder is Parkinson's disease, Alzheimer disease, or a disorder which causes some of the symptoms of Parkinson's disease.

3. The use of claim 1 or claim 2 wherein the aristolactam is Taliscanin.

4. The use of claim 3 wherein the dosage of Taliscanin to be administered is from 280 to 4000 micrograms per day, e.g. 280 micrograms per day.

5. The use of claim 3 or claim 4 wherein the Taliscanin is administered in a solution which includes ethanol and water, the alcohol concentration of the final solution preferably being between 30 and 60 proof.

6. The use of any of claims 3 to 5 wherein the medication to be administered further includes Hyaluronidase or Papavaina.

7. A solution comprising Taliscanin and a solvent which upon internal administration is effective in treating Parkinson's disease and related neurological disorders.

8. A solution comprising Taliscanin, water and dextrose, lactose or ethanol whereby upon internal administration the solution is effective in treating Parkinson's disease and related neurological disorders; and if ethanol is used the ethanol concentration optionally being 30 to 60 proof.

9. A pharmaceutical or veterinary formulation comprising an aristolactam, preferably Taliscanin, formulated for pharmaceutical or veterinary use in the treatment of neurological disorders or impotence, optionally in unit dosage form and/or optionally including an acceptable diluent, carrier or excipient.

10. Taliscanin or a solution of claim 7 or claim 8 or a formulation of claim 9 for use in either:

(a) treating Parkinson's disease and neurological disorders causing Parkinsonian symptoms by a method comprising:

administering 280 micrograms per day of Taliscanin to an afflicted patient;

increasing the dosage of Taliscanin to 600 micrograms per day if the patient does not improve; and gradually increasing the dosage of Taliscanin in increments of 280 micrograms per day until the patient improves or until the dosage reaches 4000 micrograms per day;

optionally the Taliscanin to be administered in a solution which includes ethanol or lactose or dextrose and water, and if ethanol is used preferably the alcohol concentration of the final solution is between 30 and 60 proof, and/or

optionally the method further including concurrently administering Hyaluronidase or Papavaina to the treated patient; or

(b) a method for treating Parkinson's disease comprising:

administering 280 to 4000 micrograms per day of Taliscanin to an affected patient.

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

**0 304 294
A3**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 88307636.6

(51) Int. Cl.⁵: **A61K 31/40**

(22) Date of filing: 18.08.88

(30) Priority: 20.08.87 US 87432

(43) Date of publication of application:
22.02.89 Bulletin 89/08

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(88) Date of deferred publication of the search report:
17.10.90 Bulletin 90/42

(71) Applicant: **MONOCLONETICS
INTERNATIONAL, INC.**
18333 Egret Bay Boulevard Suite 270
Houston Texas 77058(US)

(72) Inventor: **De La Parra, Jacinto Juarez**
122 Blond Petersen
Ermita Prado Mexico City MX DF(MX)

(74) Representative: **Bizley, Richard Edward et al**
BOULT, WADE & TENNANT 27 Farnival Street
London EC4A 1PQ(GB)

(54) **Taliscanin and other Aristolactams for treating neurological disorders, Parkinson's disease, Alzheimer disease and impotence.**

(57) Aristolactams e.g. Taliscanin may be used in preparing medication for treating neurological disorders, in particular Parkinson's disease, or impotence. The method involves administering an aristolactam, especially Taliscanin, an extract of the root of Aristolochia Taliscana, to an afflicted patient with a dosage varied in accordance with the patient's response, which dosage generally ranges from 280 to 4000 micrograms per day.

EP 0 304 294 A3



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT
which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

Application number

EP 88 30 7636

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	JOURNAL OF NATUREL PRODUCTS, vol. 45, no. 6, November-December 1982, pages 657-666, The American Society of Pharmacognosy and the Lloyd Library and Museum; D.B. MIX et al.: "The aristolochic acids and aristolactams" * Page 657; page 663, no. 22 * --	1-10	A 61 K 31/40
D,A	CIENCIA, vol. 24, no. 5-6, February 15, 1966, pages 237-240; L.A. MALDONADO et al.: "La taliscanina, un componente de aristolochia taliscana" * Page 237, left-hand column * --	1-10	
A	J. ORGAN. CHEM., vol. 30, November 1965, pages 3792-3800; ./. --		TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			A 61 K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-9 Claims searched incompletely: 10 Claims not searched: Reason for the limitation of the search:</p> <p>Method for treatment of the human or animal body by surgery or therapy (see Art. 52(4) of the European Patent Convention).</p>			
Place of search		Date of completion of the search	Examiner
The Hague		20-03-1990	FOERSTER
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published later, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			



PARTIAL
EUROPEAN SEARCH REPORT

Application number

EP 88 30 7636 -2-

DOCUMENTS CONSIDERED TO BE RELEVANT					
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. +)		
	S.M. KUPCHAN et al.: "Tumor inhibitors. X. Photochemical synthesis of phenanthrenes. Synthesis of aristolochic acid and related compounds"	1-10			
	* Page 3792, left-hand column; structure IV *				

			TECHNICAL FIELDS SEARCHED (Int. Cl. +)		
	XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX				
Place of search		Date of completion of the search	Examiner		
CATEGORY OF CITED DOCUMENTS			T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document		
X :	particularly relevant if taken alone				
Y :	particularly relevant if combined with another document of the same category				
A :	technological background				
O :	non-written disclosure				
P :	intermediate document				

THIS PAGE BLANK (USPTO).